

**Ambient Air Monitoring Program
Wabamun–Genesee Area**

2007 Annual Report

**Prepared for:
TransAlta Utilities Corporation and EPCOR Utilities Inc.
Edmonton, Alberta**

**Prepared by:
WBK & Associates Inc.
St. Albert, Alberta**

June 2008

Executive Summary

TransAlta Utilities Corporation and EPCOR Utilities Inc. operate four coal-fired thermal generating plants – Sundance, Keephills, Wabamun, and Genesee – located in the Wabamun–Genesee area of west-central Alberta. These generating plants operate under Alberta Environmental Protection and Enhancement Act approvals. As part of their approvals, the generating plants conduct special environmental monitoring programs, including ambient air monitoring. This 2007 annual report summarizes the results of the ambient air monitoring program that are important for understanding the state of air quality in the Wabamun–Genesee area. Components conducted during 2007 included a continuous monitoring program, an intermittent monitoring program and a passive monitoring program.

Continuous Monitoring Program

During 2007, a continuous program was conducted at the Genesee, Meadows, Wagner and Powers air monitoring stations (AMS) in the Wabamun–Genesee area. Data capture rates for measured parameters at all air monitoring stations were above the 90 percent criterion required by the Alberta Environment Air Monitoring Directive. High up-times indicated that equipment in the continuous air monitoring network was generally well-maintained. The Genesee AMS experienced a power failure in July and was down for five days, which resulted in up-times of 84% for monitoring equipment during the month. The Meadows AMS also experienced a power failure. It was down for 2.5 days in June, which resulted in up-times of 89% for monitoring equipment during the month. Both air monitoring stations operated normally for the remainder of the year.

Overall, air quality was judged to be good at the monitoring stations during 2007, similar to that observed in 2006:

- Hourly NO₂ concentrations were very low at all four air monitoring stations (<54 µg/m³ for 98 percent of the time). The observed 10th highest 1-hour NO₂ concentrations ranged from 54 µg/m³ at Genesee AMS to 70 µg/m³ at Meadows AMS. For comparison purposes, the 10th highest 1-hour NO₂ concentration observed at the Edmonton East AMS during 2007 was 115 µg/m³. All of these concentrations fall well below the 1-hour Alberta Ambient Air Quality Objective of 400 µg/m³.
- Hourly SO₂ concentrations were very low at all four air monitoring stations (<33 µg/m³ for 98 percent of the time). The observed 10th highest 1-hour SO₂ concentrations observed ranged from 72 µg/m³ at Meadows AMS to 95 µg/m³ at Powers AMS. For comparison purposes, the 10th highest 1-hour SO₂ concentration observed at the Edmonton East AMS during 2007 was 56 µg/m³. All of these concentrations fall well below the 1-hour Alberta Ambient Air Quality Objective of 450 µg/m³.
- The annual average ozone (O₃) concentration at the Genesee AMS was 53 µg/m³; similar to that observed in 2006 and 2005. Hourly O₃ concentrations at the Genesee AMS were comparable to that observed at a rural air monitoring station in west central Alberta (Carrot Creek). Hourly O₃ concentrations at the Genesee AMS were <113 µg/m³ for 98 percent of the time during 2007. The 10th highest 1-hour O₃ concentration observed was 141 µg/m³ at Genesee AMS. For comparison purposes, the 10th highest

1-hour O₃ concentrations observed at the Carrot Creek AMS and Edmonton East AMS were 139 and 135 µg/m³, respectively. All of these concentrations fall below the 1-hour Alberta Ambient Air Quality Objective of 160 µg/m³.

- Twenty-four hour PM_{2.5} concentrations at the Genesee and Powers AMS ranged from <1 to 20 µg/m³. Median (50th percentile) 24-hour PM_{2.5} concentrations at both stations were very low (≤4 µg/m³). The 98th percentile 24-hour PM_{2.5} concentrations at both air monitoring stations were low (<16 µg/m³).

Intermittent Monitoring Program

During 2007, 24-hour average particulate matter (PM₁₀ and PM_{2.5}) samples were collected at the Powers and Genesee air monitoring stations according to a National Air Pollution Surveillance six-day cycle sampling frequency:

- Twenty-four hour average PM_{2.5} concentrations were <20 µg/m³ at both the Genesee AMS and Powers AMS for 98 percent of the time. This is consistent with 24-hour average data observed from continuous monitoring at these stations.
- Twenty-four hour average PM₁₀ concentrations were <51 µg/m³ at both the Genesee and Powers air monitoring stations for 98 percent of the time.

Passive Monitoring Program

Passive air monitors were deployed at 21 sites in the Wabamun–Genesee area during 2007:

- Annual average NO₂ concentrations ranged from 4.5 to 7.3 µg/m³ at passive sites located within 30 km of the generating plants, whereas the annual average NO₂ concentration observed at the Edmonton East AMS was 29 µg/m³. Annual NO₂ concentrations increased slightly from sites near Lake Wabamun to sites adjacent to the City of Edmonton.

This spatial trend is consistent with previous studies in the same area. It is due to the influence of increasing urban NO_x emissions within and adjacent the City of Edmonton. Urban source emissions are more important contributors to ambient NO₂ concentrations observed in the City of Edmonton and surrounding area, compared with generating plant emissions in the Wabamun–Genesee area.

- Annual average SO₂ concentrations were very low – in the 2.6 to 5.2 µg/m³ range – at all passive sites in the Wabamun–Genesee area. These low concentrations and associated lack of spatial trend is consistent with that reported in previous studies for the same area.
- At passive monitoring sites located directly east of the Wabamun–Genesee area, the observed annual average O₃ concentrations ranged from 23 to 29 µg/m³; whereas at the Violet Grove AMS, a rural background air monitoring station located 55 km west of the generating plants, an annual average O₃ concentration of 29 µg/m³ was observed.
- Two passive monitoring sites are located greater than two- to four-hours travel time downwind of the generating plants along an imaginary northwest line on which the generating plants are located. Previous work by others has indicated that generating

plant plume chemistry maturity and peak production capacities of anthropogenic O₃ and inorganic nitrogen species occurs between 30 and 100 km downwind of generating plants (within the range of where these sites are located).

Annual average O₃ concentrations at these two sites were within five percent of the annual average O₃ concentration at the Violet Grove AMS. This difference is within the expected accuracy and precision of the O₃ passive monitoring device used (± 15 percent). These findings indicate that observation of anthropogenic O₃ production downwind of generating plant emissions is not readily apparent using passive monitors.

Table of Contents

Executive Summary	i
Table of Contents.....	iv
List of Figures	v
List of Tables.....	vi
Abbreviations	vii
1 Introduction	1-1
1.1 Environmental Monitoring Programs for Generating Plants	1-2
1.2 Purpose of Report	1-4
2 Background on Air Quality Parameters Monitored.....	2-1
2.1 Oxides of Nitrogen.....	2-1
2.2 Sulphur Dioxide	2-1
2.3 Ground-Level Ozone	2-1
2.4 Particulate Matter	2-2
3 Methods	3-1
3.1 Continuous Monitoring Program.....	3-1
3.2 Intermittent Monitoring Program.....	3-2
3.3 Passive Monitoring Program	3-2
4 Results and Discussion.....	4-1
4.1 Continuous Monitoring Program.....	4-1
4.1.1 Percent Completeness	4-1
4.1.2 Air Pollutant Concentration Statistics	4-2
4.1.3 Wind Speed and Direction Statistics	4-11
4.2 Intermittent Monitoring Program.....	4-16
4.3 Passive Monitoring Program	4-18
4.3.1 Nitrogen Dioxide	4-19
4.3.2 Sulphur Dioxide	4-19
4.3.3 Ozone	4-19
5 Findings.....	5-1
6 References.....	6-1
APPENDIX Summary of 2007 Passive Monitoring Data.....	A-1

List of Figures

Figure 1-1	Locations of Coal-fired Generating Plants and Air Monitoring area for Generating Plants in the Wabamun-Genesee Area.	1-1
Figure 1-2	Continuous and Passive Monitoring Locations in the Wabamun-Genesee Area during 2007	1-3
Figure 4-1	Box-and-Whisker Plot of 1-hour Average NO ₂ Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007.....	4-3
Figure 4-2	Box-and-Whisker Plot of 24-hour Average NO ₂ Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007.....	4-4
Figure 4-3	Cumulative Distribution Plot of 1-hour Average NO ₂ Concentrations at Air Monitoring Stations in the Wabamun-Genesee Area during 2007	4-5
Figure 4-4	Box-and-Whisker Plot of 1-hour Average SO ₂ Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007.....	4-6
Figure 4-5	Box-and-Whisker Plot of 24-hour Average SO ₂ Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007.....	4-7
Figure 4-6	Cumulative Distribution Plot of 1-hour Average SO ₂ Concentrations at Air monitoring Stations in the Wabamun-Genesee Area during 2007	4-8
Figure 4-7	Box-and-Whisker Plot of 1-hour Average O ₃ Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007.....	4-8
Figure 4-8	Cumulative Distribution Plot of 1-hour Average O ₃ Concentrations at the Genesee Air Monitoring Station during 2007.....	4-9
Figure 4-9	Box-and-Whisker Plot of 24-hour Average PM _{2.5} Concentrations at the Genesee and Powers Air Monitoring Stations (2005, 2006, and 2007).....	4-10
Figure 4-10	Cumulative Distribution Plot of 1-hour Average PM _{2.5} Concentrations at the Powers and Genesee Air Monitoring Stations during 2007.....	4-11
Figure 4-11	Annual Wind Rose Plot for the Genesee Air Monitoring Station during 2007.....	4-12
Figure 4-12	Annual Wind Class Frequency Distribution Plot for the Genesee Air Monitoring Station during 2007.....	4-13
Figure 4-13	Annual Wind Rose Plot for the Powers Air Monitoring Station during 2007.....	4-13
Figure 4-14	Annual Wind Class Frequency Distribution Plot for the Powers Air Monitoring Station during 2007.....	4-14
Figure 4-15	Annual Wind Rose Plot for the Meadows Air Monitoring Station during 2007.....	4-14
Figure 4-16	Annual Wind Class Frequency Distribution Plot for the Meadows Air Monitoring Station during 2007.....	4-15
Figure 4-17	Annual Wind Rose Plot for the Wagner Air Monitoring Station during 2007.....	4-15
Figure 4-18	Annual Wind Class Frequency Distribution Plot for the Wagner Air Monitoring Station during 2007.....	4-16
Figure 4-19	Box-and-Whisker Plot of Intermittent 24-hour Average PM _{2.5} Concentrations at Genesee and Powers Air Monitoring Stations in 2007.....	4-16

Figure 4-20	Box-and-Whisker Plot of 24-hour Average PM ₁₀ Concentrations at Genesee and Powers Air Monitoring Stations in 2007	4-17
Figure 4-21	Twenty-four Hour Average PM _{2.5} Concentration versus PM ₁₀ Concentration at the Genesee Air Monitoring Station during 2007	4-17
Figure 4-22	Twenty-four Hour Average PM _{2.5} Concentration versus PM ₁₀ Concentration at the Powers Air Monitoring Station during 2007	4-18
Figure 4-23	Bubble Plot of Annual Average NO ₂ Concentrations [ppb] at Passive Monitoring Sites and at Selected Continuous Monitoring Station Locations in Central Alberta during 2007	4-20
Figure 4-24	Bubble Plot of Annual Average SO ₂ Concentrations [ppb] at Passive Monitoring Sites and at Selected Continuous Monitoring Station Locations in Central Alberta during 2007	4-21
Figure 4-25	Bubble Plot of Annual Average O ₃ Concentrations [ppb] at Passive Monitoring Sites and at Selected Continuous Monitoring Station Locations in Central Alberta during 2007	4-22
Figure 4-26	Monthly Average O ₃ Concentration at Passive Monitoring Sites in Central Alberta during 2007	4-24

List of Tables

Table 1-1	Alberta Environmental Protection and Enhancement Act (EPEA) Operating Approvals for Four Generating Plants in the Wabamun-Genesee Area.....	1-2
Table 2-1	Air Quality Objectives and Standards for Specific COPCs Related to Generating Plant Emissions in the Wabamun-Genesee Area.....	2-3
Table 3-1	Sampling Schedule for Parameters in the Ambient Air Quality Monitoring Program in the Wabamun-Genesee Area	3-3
Table 4-1	Monthly and Annual Data Capture Rates (% completeness) for Genesee Air Monitoring Station during 2007	4-1
Table 4-2	Monthly and Annual Data Capture Rates (% completeness) for Meadows Air Monitoring Station during 2007	4-1
Table 4-3	Monthly and Annual Data Capture Rates (% completeness) for Wagner Air Monitoring Station during 2007	4-2
Table 4-4	Monthly and Annual Data Capture Rates (% completeness) for Powers Air Monitoring Station during 2007	4-2

Abbreviations

AAAQO	Alberta ambient air quality objective
AENV	Alberta Environment
AMS	air monitoring station
CASA	Clean Air Strategic Alliance
COPCs	Chemicals of Potential Concern
CWS	Canada-Wide Standard
EPEA	Environmental Protection and Enhancement Act
IQR	interquartile range
NAPS	National Ambient Pollutant Surveillance
NO ₂	nitrogen dioxide
NO _x	oxides of nitrogen
O ₃	ozone
ppb	parts per billion
PM _{2.5}	particulate matter ≤2.5 µm
PM ₁₀	particulate matter ≤10 µm
R ²	coefficient of determination
SO ₂	sulphur dioxide
TSP	total suspended particulate
MW	megawatts
U.S. EPA	United States Environmental Protection Agency
VOCs	volatile organic compounds
WCAS	West Central Airshed Society

1 Introduction

TransAlta Utilities Corporation (TransAlta) [www.transalta.com] and EPCOR Utilities Inc. (EPCOR) [www.epcor.ca] operate four coal-fired thermal generating plants (generating stations) Wabamun–Genesee area of west-central Alberta. The generating plants include Wabamun, Sundance, Keephills, and Genesee. The location of these plants is shown in Figure 1-1. Collectively, the four generating plants have a net generating capacity of 4277 MW.




Figure 1-1 Locations of Coal-fired Generating Plants and Air Monitoring Area for Generating Plants in the Wabamun–Genesee Area

The Wabamun generating plant is the oldest of TransAlta’s three generating plants in the Lake Wabamun area. It is near the Village of Wabamun and has a net generating capacity of 279 MW (Figure 1-1). Only one generating unit was in operation at the Wabamun plant in 2007. The remaining three units were retired in 2002 (Unit 3) and 2004 (Units 1 and 2).

The TransAlta Sundance generating plant consists of six generating units. It is the largest, coal-fired generating plant in western Canada. Sundance is situated on the south shore of Lake Wabamun, approximately 70 km west of Edmonton, Alberta (Figure 1-1).

The plant has been in operation since 1970, with steady expansion from a single original generating unit to six generating units throughout the 1970s. Sundance currently has a net generating capacity of 2020 MW.

The Keephills generating plant is located 5 km southeast of Lake Wabamun (Figure 1-1). It has a net generating capacity of 766 MW, and consists of two generating units. Keephills has been in operation since 1983.

The Genesee generating plant, located 50 km southwest of Edmonton, consists of three generating units (Figure 1-1). EPCOR fully owns and operates Units 1 and 2, which have a combined net generating capacity of 762 MW. These units have been in operation since 1994 and 1989, respectively. Genesee 3 (Unit 3), commissioned in 2005, is a 50/50 joint venture between TransAlta and EPCOR. Genesee 3 has a net generating capacity of 450 MW.

1.1 Environmental Monitoring Programs for Generating Plants

The generating plants operate under the Alberta Environmental Protection and Enhancement Act (EPEA) approvals listed in Table 1-1. Under their EPEA approvals, the generating plants are committed to conducting special environmental monitoring programs. These programs are designed to:

- Identify and quantify ambient levels and deposition patterns of chemical species of potential concern that are associated with generating plant emissions.
- Generate an inventory of representative baseline data for the chemicals of potential concern.
- Provide data for assessing long-term impacts and for evaluating and implementing air quality management plans.

Table 1-1 Alberta Environmental Protection and Enhancement Act (EPEA) Operating Approvals for Four Generating Plants in the Wabamun-Genesee Area

Facility	Capacity (MW, net)	Location	Approval No. (as amended)	Applicable Approval Terms
Wabamun	279	2,3,10,11-53-04 W5M	10323-02-00	6.1.14 to 6.1.24; 6.1.32 to 6.1.34
Sundance	2020	3,4,8,9,10,16,17,20, and 31-52-04 W5M	9830-02-00	7.1.1 to 7.1.5
Keephills	766	36-51-04 W5M	10324-01-00	6.1.14 to 6.1.24; 6.1.32 to 6.1.34
Genesee	1212	25-50-03 W5M	773-02-00	7.1.1 to 7.1.5




Figure 1-2 Continuous and Passive Monitoring Locations in the Wabamun-Genesee Area during 2007

A component of the special environmental monitoring program, includes developing and implementing an ambient air quality monitoring program. The ambient air quality monitoring program consists of the following elements (Figure 1-2):

- A continuous monitoring program for: sulphur dioxide (SO₂), nitrogen dioxide (NO₂), and a number of meteorological parameters at four air monitoring stations (AMS) (Genesee AMS, Powers AMS, Meadows AMS, and Wagner AMS); particulate matter with aerodynamic diameter less than or equal to 2.5 µm (PM_{2.5}) at two stations (Genesee AMS and Powers AMS); and ozone (O₃) at one station (Genesee AMS).
- An intermittent monitoring program (24 hour sampling every 6 days) for particulate matter (with aerodynamic diameter less than or equal to 10 µm PM₁₀, and PM_{2.5}, and metals speciation of the PM_{2.5} at two stations (Genesee AMS and Powers AMS).
- A passive monitoring program consisting of monthly passive monitoring for NO₂, SO₂, and O₃ at 21 locations in the Wabamun–Genesee area..

Specific clauses within the EPEA approvals that refer to the ambient air quality monitoring program are provided in Table 1-1.

1.2 Purpose of Report

This annual report analyzes and presents information collected as part of the 2007 ambient air monitoring program. These data provide a basis for developing an understanding of the state of air quality in relation to emissions from the four coal-fired generating plants in the Wabamun–Genesee area. Monitoring components associated with an acid deposition assessment program and a mercury assessment program in the Wabamun–Genesee area are reported separately in stand-alone documents (TransAlta and EPCOR 2008a, b).

2 Background on Air Quality Parameters Monitored

A number of chemicals of potential concern (COPCs) are potentially emitted from the coal-fired generating plants in the Wabamun–Genesee area. With respect to the ambient air monitoring program, these chemicals of potential concern include: oxides of nitrogen, sulphur dioxide, ground level ozone, and airborne particulate matter. The importance of these COPCs is discussed below.

2.1 Oxides of Nitrogen

Oxides of nitrogen (NO_x) are made up of nitrogen dioxide (NO_2) and nitric oxide (NO). They are formed during high temperature combustion processes when nitrogen present in both hydrocarbon fuel and combustion air react with oxygen in combustion air to form NO. NO quickly reacts with ozone in ambient air to form NO_2 (Environment Canada 2006a). The major emission sources of NO_x are transportation, electrical utilities, and industrial, commercial and residential fuel-burning sources (U.S. EPA 1998a). NO_x contributes to formation of ozone, acid aerosols, nitrate particulates, acid rain, and nutrient overload that result in lower water quality (U.S. EPA 1998a).

The Alberta ambient air quality objective (AAAQO) for NO_x is expressed as NO_2 . NO_x concentrations are typically expressed as NO_2 for comparison to an AAAQO. The Clean Air Strategic Alliance (CASA) (2006) reports that a 10-year average concentration of NO_2 measured at a rural background location in Alberta (Hightower Ridge) is 8 ppb ($15 \mu\text{g}/\text{m}^3$).

2.2 Sulphur Dioxide

Sulphur dioxide (SO_2) is formed from combustion of sulphur containing raw materials such as coal, fuel oils, and metallic ores (Environment Canada 2006b). The SO_2 reacts with water vapour in air to form acidic sulphur compounds and particulates (U.S. EPA 2000). Major sources of SO_2 are metal smelters, fossil fuel-fired generating plants, transportation, upstream oil and gas, and other industrial facilities. CASA (2006) reports that a 10-year average concentration of SO_2 measured at a rural background location in Alberta (Hightower Ridge) is 2 ppb ($5 \mu\text{g}/\text{m}^3$).

2.3 Ground-Level Ozone

Ground-level ozone (O_3) is a secondary pollutant, which means that it is not directly emitted into the atmosphere. It is formed through a series of photochemical reactions between NO_x and volatile organic compounds (VOCs). Major sources of NO_x were described above. Major emission sources of VOCs are both natural (vegetation) and anthropogenic (motor vehicles, petroleum and chemical industries, and other combustion processes). Ground-level ozone may also occur through intrusion of upper stratospheric ozone into the troposphere (ground-level layer of the atmosphere).

CASA (2006) reports that a 10-year average concentration of O₃ measured at a rural background location in Alberta (Hightower Ridge) is 42 ppb (81 µg/m³).

He et al. (2005) and Kindzierski et al. (2005) reported that meteorological conditions appear to be the most important factors related to the behaviour of O₃ concentrations (e.g., hourly average O₃ concentration maxima and minima) observed at three air monitoring stations in west-central Alberta.

It was found by these authors that variation in hourly O₃ concentrations was closely related to changes in hourly temperature, relative humidity, and pressure. This strongly points to the importance of natural phenomena as the predominant factor contributing to the presence of ground-level O₃ at these monitoring sites. Anthropogenic factors (i.e., precursor air pollutants originating from the activity of humans) were less important to the behaviour of hourly O₃ concentrations at these monitoring sites.

2.4 Particulate Matter

Airborne particulate matter (PM) is a complex mix of various pollutants in solid and liquid forms in ambient air. These pollutants include acids (sulphates and nitrates), metals, organic chemicals, and soil or dust particles (U.S. EPA 2006). Airborne particulate matter is commonly distinguished by three different size fractions:

- Total suspended particulate (TSP) – particles with aerodynamic diameters 100 µm or less.
- PM₁₀ – particles with aerodynamic diameters 10 µm or less.
- PM_{2.5} – particles with aerodynamic diameters 2.5 µm or less.

CASA (2006) reports that a 10-year average concentration of PM_{2.5} measured at a rural background location (Hightower Ridge) is 2.3 µg/m³.

2.5 Ambient Air Objectives

Air quality criteria are used to protect human health, safeguard the environment, and for aesthetic purposes. In this case, specific Alberta Ambient Air Quality Objectives (AAAQO) and Canada-wide Standards (CWS) exist for these COPCs to aid in the interpretation of measured ambient levels of these pollutants. These objectives and standards are listed in Table 2-1.

Table 2-1 Air Quality Objectives and Standards for Specific COPCs Related to Generating Plant Emissions in the Wabamun–Genesee Area

Parameter	Averaging Time	Alberta Ambient Air Quality Objective	Canada-Wide Standard
NO _x (as NO ₂)	1 hour	400 µg/m ³ (212 ppb)	-
	24 hour	200 µg/m ³ (106 ppb)	-
	annual	60 µg/m ³ (32 ppb)	-
SO ₂	1 hour	450 µg/m ³ (172 ppb)	-
	24 hour	150 µg/m ³ (57 ppb)	-
	annual	30 µg/m ³ (11 ppb)	-
PM _{2.5}	24-hour average based on 98 th percentile value over 3 consecutive years	-	30 µg/m ³
O ₃	1 hour	160 µg/m ³ (82 ppb)	-
	8 hour average based on 4 th highest value over 3 consecutive years	-	128 µg/m ³ (65 ppb)

NOTE:
 - not applicable

3 Methods

3.1 Continuous Monitoring Program

Analytical methods used to determine ambient pollutant concentrations for the continuous program conducted at the Genesee, Meadows, Wagner, and Powers air monitoring stations are described in the following sections. Refer to Figure 1-2 for monitoring locations. Procedures and guidelines for measuring and analyzing the air quality parameters listed below are described further in the Ambient Air Monitoring Program Quality Assurance Plan for West Central Airshed Society (WBK 2007).

NO₂

NO₂ is analyzed at Genesee, Meadows, Wagner, and Powers air monitoring stations. NO₂ is measured by chemiluminescence using TECO 42C (Thermo Electron Corporation, Waltham, MA) or equivalent analyzers. NO_x in air is converted to nitric oxide (NO) as it flows over a heated catalyst. The NO is then oxidized by O₃, resulting in light emission during the reaction. The light emission, which is proportional to each molecule of NO that is oxidized, is measured and presented on an NO₂ basis.

SO₂

SO₂ is analyzed at Genesee, Meadows, Wagner, and Powers air monitoring stations. SO₂ is measured with pulsed fluorescence using TECO 43 (Thermo Electron Corporation, Waltham, MA) or equivalent analyzers. Air samples are bombarded with pulses of ultraviolet (UV) light that excite SO₂ molecules to higher energy levels. When the SO₂ molecules return to their original energy state, light is emitted; this light is measured in the analyzer and is proportional to the amount of SO₂ in air.

O₃

O₃ is measured at the Genesee AMS with a TECO 49 UV-absorption ozone analyzer (Thermo Electron Corporation, Waltham, MA). This measurement is based on ozone's ability to absorb UV light. A sample of air is exposed to UV light and the amount absorbed is measured. The UV light absorbed is proportional to the amount of O₃ in air.

PM_{2.5}

PM_{2.5} is measured at Powers AMS and Genesee AMS using a tapered element oscillating microbalance (TEOM) (Thermo Electron Corporation, Waltham, MA). Particulates in air are separated by diameter and passed through a filter attached to a tapered element vibrating at its natural frequency. As particles deposit on the filter, the vibration frequency of the element changes in proportion to the mass of particles deposited.

Meteorology

Wind speed, wind direction, air temperature, and relative humidity are monitored at Genesee, Meadows, Wagner, and Powers air monitoring stations using standard meteorological monitoring equipment.

In 2007, West Central Airshed Society (WCAS) technicians maintained and calibrated the analyzers at the continuous stations. The data are reviewed for errors and omissions by WCAS personnel and the information is reported on a monthly, quarterly, and annual basis. The raw data are also available to the public on the WCAS website (www.wcas.ca) on a real-time basis.

3.2 Intermittent Monitoring Program

Twenty-four hour average PM₁₀ and PM_{2.5} samples are collected at Powers and Genesee air monitoring stations. These samples are collected according to a National Air Pollution Surveillance (NAPS) six-day cycle sampling frequency using Partisols (Thermo Electron Corporation, Waltham, MA). Procedures and guidelines for measuring and analyzing PM₁₀ and PM_{2.5} samples are described further in the Ambient Air Monitoring Program Quality Assurance Plan for West Central Airshed Society (WBK 2007).

The Partisols separate particulates in sampled air according to size, which allows for size-selective collection on a filter. Technicians from Ambitech Inc. (Edmonton, AB) collected Partisol filters according to the NAPS schedule (see http://www.etc-cte.ec.gc.ca/NAPS/index_e.html) and shipped the filters to the Alberta Research Council (Vegreville, AB) for gravimetric and metals analyses. WBK & Associates Inc. personnel reviewed the analytical results on an as-received basis.

Results from metals analyses of 24-hour average PM_{2.5} samples are currently archived electronically. They will eventually be used to undertake source apportionment, which is the quantitative identification of the relative contributions of different source types to airborne particle concentrations. Source apportionment for PM_{2.5} will define the total contribution that different sources in the Wabamun–Genesee area make to airborne particle concentrations at the Powers and Genesee air monitoring stations. Generally, greater than 100 PM_{2.5} samples are required to undertake source apportionment (≥2 years of monitoring data).

3.3 Passive Monitoring Program

Passive air monitors are deployed at a total of 21 stations in the Wabamun–Genesee area (Figure 1-2). Maxxam Analytics (Edmonton, AB) PASS samplers are deployed for durations of approximately one month to measure SO₂, NO₂, and O₃. Procedures and guidelines for collecting and analyzing SO₂, NO₂, and O₃ passive samplers are described further in the Ambient Air Monitoring Program Quality Assurance Plan for West Central Airshed Society (WBK 2007).

The passive approach collects gas from the atmosphere at a rate controlled by the gas' natural diffusion across a membrane. Passive collection of a given air pollutant is achieved by chemical absorption or by physical adsorption onto a medium in the sampler. Ambitech Inc. technicians collected the passive samplers and deployed fresh samplers within two days of the end of each month. The samplers were delivered to Maxxam Analytics (Edmonton, AB) for laboratory analysis.

A schedule for the sampling programs described above is presented in Table 3-1.

Table 3-1 Sampling Schedule for Parameters in the Ambient Air Quality Monitoring Program in the Wabamun–Genesee Area

Parameter	Continuous	Intermittent (every 6 th day (NAPS) schedule)	Monthly (passives)
SO ₂	•		•
NO ₂	•		•
O ₃	•		•
PM ₁₀		•	
PM _{2.5}	•	•	
Wind speed and direction, temperature, relative humidity)	•		

4 Results and Discussion

4.1 Continuous Monitoring Program

4.1.1 Percent Completeness

An important component for ensuring quality for continuous monitoring data is completeness. Completeness indicates the percentage of time that a continuous monitor is capturing valid data. In general, the Air Monitoring Directive (1989) requires greater than 90 percent completeness. Tables 4-1 to 4-4 show data capture rates (percent completeness) for the four continuous air monitoring stations in the Wabamun–Genesee area during 2007.

Table 4-1 Monthly and Annual Data Capture Rates (% completeness) for Genesee Air Monitoring Station during 2007

Parameter	Up-time (%)												Annual
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
NO ₂	100	100	100	100	99.6	99.7	83.7	99.9	99.9	99.3	100	100	98.5
SO ₂	100	100	100	100	99.6	99.7	83.7	99.9	99.4	98.5	100	100	98.4
O ₃	100	100	100	100	99.6	99.7	83.7	99.9	99.9	99.2	100	100	98.5
PM _{2.5}	99.6	99.3	100	99.3	99.6	99.8	81.6	99.1	99.4	98.3	100	97.8	97.8
WSP	100	100	100	100	99.6	99.7	83.7	99.9	99.9	99.3	100	96.6	98.2
WDR	100	100	100	100	99.6	99.7	83.7	99.9	99.9	99.3	100	96.6	98.2
T ₂	100	100	100	100	99.6	99.7	83.7	99.9	99.7	99.3	100	100	98.5
T ₁₀	100	100	100	100	99.6	99.7	83.7	99.9	99.9	99.3	100	100	98.5
RH	100	100	100	100	99.6	99.7	83.7	99.9	99.9	99.3	100	100	98.5

NOTES:

Key: WSP = wind speed; WDR = wind direction; T₂ = temperature at 2-metre height above ground; T₁₀ = temperature at 10-metre height above ground; RH = relative humidity.

Table 4-2 Monthly and Annual Data Capture Rates (% completeness) for Meadows Air Monitoring Station during 2007

Parameter	Up-time (%)												Annual
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
NO ₂	98.5	99.6	99.9	98.6	98.4	88.8	98.5	99.5	100	99.9	99.4	99.9	98.4
SO ₂	98.5	100	99.9	98.6	98.4	88.8	98.5	99.5	100	99.9	99.4	99.9	98.5
WSP	98.5	100	99.9	98.6	98.4	88.8	99.9	99.5	100	99.9	99.4	99.9	98.6
WDR	98.5	100	99.9	98.6	98.4	88.8	99.9	99.5	100	99.9	99.4	99.9	98.6
T ₂	98.5	100	99.9	98.6	98.4	88.8	99.9	99.5	100	99.9	99.4	99.9	98.6
RH	98.5	100	99.9	98.6	98.4	88.8	99.9	99.5	100	99.9	99.4	99.9	98.6

NOTES:

Key: WSP = wind speed; WDR = wind direction; T₂ = temperature at 2-metre height above ground; RH = relative humidity.

Table 4-3 Monthly and Annual Data Capture Rates (% completeness) for Wagner Air Monitoring Station during 2007

Parameter	Up-time (%)												Annual
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
NO ₂	99.9	100	100	100	100	96.7	100	100	100	100	100	99.7	99.7
SO ₂	99.9	100	100	100	100	96.7	100	100	100	100	100	99.7	99.7
WSP	99.9	100	100	100	100	96.8	100	100	100	100	100	100	99.7
WDR	99.9	100	100	100	100	96.8	100	100	100	100	100	100	99.7
T ₂	99.9	100	100	100	100	96.8	100	100	99.9	100	100	100	99.7
RH	99.9	100	100	100	100	96.8	100	100	100	100	100	100	99.7

NOTES:

Key: WSP = wind speed; WDR = wind direction; T₂ = temperature at 2-metre height above ground; RH = relative humidity.

Table 4-4 Monthly and Annual Data Capture Rates (% completeness) for Powers Air Monitoring Station during 2007

Parameter	Up-time (%)												Annual
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
NO ₂	100	100	100	100	100	100	99.9	100	100	100	100	100	100
SO ₂	100	100	100	100	100	100	99.9	100	100	100	100	100	100
PM _{2.5}	99.7	100	98.7	99.9	99.6	98.5	96.1	98.4	97.9	99.9	99.4	100	99.0
WSP	100	100	100	100	100	100	99.9	100	100	100	100	100	100
WDR	100	100	100	100	100	100	99.9	100	100	100	100	100	100
T ₂	100	100	100	100	100	100	99.9	100	99.7	100	100	100	100
RH	100	100	100	100	100	100	99.9	100	99.7	100	100	100	100

NOTES:

Key: WSP = wind speed; WDR = wind direction; T₂ = temperature at 2-metre height above ground; RH = relative humidity.

Monthly data capture rates for all stations were well above the 90 percent criterion stipulated in the Air Monitoring Directive (1989), except as noted. High up-times indicate that equipment in the continuous air monitoring network was generally well-maintained. The Genesee AMS experienced a power failure on July 17th which damaged electrical circuitry. The station was not brought back online until July 22nd, resulting in up-times of 83.7%. This incident was reported to Alberta Environment on July 20th (Reference Number: 190305). In addition, the PM_{2.5} analyzer experienced unstable operation yielding an up-time of 81.6% in July. The Meadows AMS experienced power failure spanning June 11th at 1730 hrs through June 14th at 1950 hrs, returning up-times for all analyzers and meteorological equipment of 88.8%. This event was reported to Alberta Environment on June 15th (Reference Number: 188059).

4.1.2 Air Pollutant Concentration Statistics

One method of displaying a set of air quality data is with box-and-whisker plots. Box-and-whisker plots are helpful in interpreting the distribution of data because they only illustrate certain statistics rather than all of the data. The plots presented here show five values for individual pollutants collected at each air monitoring station during 2007: 25th percentile (bottom of box), 50th percentile (line inside box), 75th percentile (top of box), maximum (top T), and annual arithmetic mean concentration (star symbol). In some cases, the 98th percentile (diamond symbol) rather than the annual arithmetic mean concentration is shown. The bottom

whisker is not shown in these plots because the values represented by bottom whiskers are not essential for data interpretation.

Nitrogen Dioxide – Figures 4-1 and 4-2 are box-and-whisker plots that compare key statistical values for 1-hour and 24-hour NO₂ concentrations observed at: the four ambient air monitoring stations, a background station relative to the air monitoring area for generating plants (Carrot Creek AMS after CASA 2008), and an urban station (Edmonton East AMS after CASA 2008). In Figures 4-1 and 4-2, the bottom of the yellow zone, boundary of the yellow and blue zones, top of the blue zone, and top tee represent the 25th, 50th, 75th, and maximum values, respectively. The star symbol in each plot represents the annual arithmetic mean concentration.




Figure 4-1 Box-and-Whisker Plot of 1-hour Average NO₂ Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007

Note: 25th %ile (bottom of box), 50th %ile (line inside box), 75th %ile (top of box), maximum (top T), and annual arithmetic mean concentration (star symbol).

Figures 4-1 (1-hour) and 4-2 (24-hour) show that NO₂ concentrations at all stations were well below the respective AAAQO for both 1-hour and 24-hour averaged concentrations. Concentration maxima – such as the maximum 1-hour concentration for a year – are typically associated with rare meteorological events (AENV 2000). These maxima can occur under conditions where emissions from sources within reasonable proximity to where monitoring is conducted (e.g., within 5 km) are fairly constant over the course of the year. For discussion purposes, the 10th highest 1-hour concentration observed (\approx 99.9th annual percentile) during 2007 is presented for the air monitoring stations to illustrate concentration maxima that are relatively free of rare meteorological influences. The 10th highest 1-hour NO₂ concentrations observed ranged from 54 µg/m³ at Genesee AMS to 70 µg/m³ at Meadows AMS during 2007.

For comparison purposes, the 10th highest 1-hour NO₂ concentrations observed at the Carrot Creek AMS and Edmonton East AMS were 72 and 115 µg/m³, respectively (CASA 2008).




Figure 4-2 Box-and-Whisker Plot of 24-hour Average NO₂ Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007

Note: 25th %ile (bottom of box), 50th %ile (line inside box), 75th %ile (top of box), maximum (top T), and annual arithmetic mean concentration (star symbol).

NO₂ concentrations were highest at the urban (Edmonton East) AMS for both 1-hour and 24-hour averaging periods, reflecting the influence of motor vehicles emissions on ambient concentrations at this urban location. Annual average NO₂ concentrations ranged from 6 µg/m³ at Powers AMS to 12 µg/m³ at Meadows AMS. These concentrations are only slightly above the 10-year average concentration reported by CASA (2006) for Hightower Ridge AMS (5 µg/m³). There were no exceedances of the AAAQO for NO₂ in 2007.

The interquartile range (IQR) is the distance between the 75th percentile and the 25th percentile for a set of data (i.e., height of each box in Figures 4-1 and 4-2). The IQR provides a good quantitative indication of the variation or degree to which values in a set of data are spread out or clustered together, because it is not affected by outliers or extreme values. Of the four air monitoring stations, the largest IQR (i.e., largest box) occurred for the Meadows AMS for hourly data (Figure 4-1) and for 24-hour data (Figure 4-2). However, IQR for NO₂ at the Meadows AMS was narrow (e.g., only 16 µg/m³ for hourly concentrations), indicating little variation in hourly concentrations throughout 2007.

Another method of displaying a set of air quality data is with a cumulative distribution plot. A cumulative distribution plot shows the fraction (or percentage) of concentration values for a pollutant that is less than or equal to a particular value. Figure 4-3 shows a cumulative

distribution plot for hourly NO₂ concentrations at each of the stations in the Wabamun–Genesee area.

Key percentile values (50th, 65th, 80th, 95th, and 98th) are shown in this plot to assist in comparison with future data and allow examination of trends in ambient air pollutant concentrations using simple methods described by Faisal et al. (2006), Haque et al. (2006) and Xu et al. (2006).

These key percentile values can be calculated from year to year to examine changes to air quality over time. Statistically significant changes in these key percentile values over a period of several years (e.g., >5 years) are indicative of trends (changes) in the concentration data. Cumulative distribution data from the Genesee, Powers and Wagner air monitoring stations (Figure 4-3) followed the same general distribution. The Meadows AMS data exhibited a greater concentration variation between the 50th and 98th percentile values compared with the other stations. Overall, hourly NO₂ concentrations were low at all four air monitoring stations (<55 µg/m³ for 98 percent of time during 2007).




Figure 4-3 Cumulative Distribution Plot of 1-hour Average NO₂ Concentrations at Air Monitoring Stations in the Wabamun–Genesee Area during 2007

Sulphur Dioxide – Figures 4-4 and 4-5 are box-and-whisker plots for 1-hour and 24-hour SO₂ concentrations, respectively. As shown in Figure 4-4, median (50th percentile) hourly SO₂ concentrations at all four generating plant air monitoring stations were low (1 µg/m³). All 1-hour and 24-hour concentrations were well below AAQOs. The 10th highest 1-hour SO₂ concentrations observed during 2007, ranged from 72 µg/m³ at Meadows AMS to 95 µg/m³ at Powers AMS. For comparison purposes, the 10th highest 1-hour SO₂ concentrations observed

at the Carrot Creek AMS and Edmonton East AMS were 33 and 56 $\mu\text{g}/\text{m}^3$, respectively (after CASA 2008).

Annual average SO_2 concentrations at the four air monitoring stations represented approximately one percent of the hourly AAAQO and two percent of the 24-hour AAAQO for SO_2 . Annual mean 1-hour and 24-hour concentrations were less than the 10-year average measured at Hightower Ridge (5 $\mu\text{g}/\text{m}^3$). There were no exceedances of any AAAQO in 2007.

Of the air monitoring stations, the Wagner AMS had the largest IQR (i.e., largest box) for hourly data (Figure 4-4) and for 24-hour data (Figure 4-5). However, IQRs for SO_2 at the Wagner AMS are very narrow (e.g., only 3 $\mu\text{g}/\text{m}^3$ for hourly concentrations), indicating extremely small variation in hourly concentrations throughout 2007.

Figure 4-6 shows the cumulative distribution and key percentile values for hourly SO_2 concentrations at the Genesee, Wabamun, Meadows and Powers air monitoring stations. SO_2 cumulative distribution curves were similar at all four air monitoring stations. Overall, hourly SO_2 concentrations were very low at all four stations (<33 $\mu\text{g}/\text{m}^3$ for 98 percent of time during 2007).




Figure 4-4 Box-and-Whisker Plot of 1-hour Average SO_2 Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007

Note: 25th %ile (bottom of box), 50th %ile (line inside box), 75th %ile (top of box), maximum (top T), and annual arithmetic mean concentration (star symbol).




Figure 4-5 Box-and-Whisker Plot of 24-hour Average SO₂ Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007

Note: 25th %ile (bottom of box), 50th %ile (line inside box), 75th %ile (top of box), maximum (top T), and annual arithmetic mean concentration (star symbol).

Ozone – Figure 4-7 is a box-and-whisker plot of 1-hour O₃ concentrations measured at the Genesee, Carrot Creek and Edmonton East air monitoring stations. The annual hourly average O₃ concentration at the Genesee AMS was 51 µg/m³, which is similar to the annual hourly average O₃ concentration observed in 2006 (53 µg/m³). The 10th highest 1-hour O₃ concentration observed in 2007 at Genesee AMS was 141 µg/m³. For comparison purposes, the 10th highest 1-hour O₃ concentrations observed at the Carrot Creek AMS and Edmonton East AMS were 139 and 135 µg/m³, respectively (after CASA 2008). The smallest IQR for O₃ (27 µg/m³) occurred at the Genesee AMS for hourly data, indicating that hourly O₃ concentrations showed the least amount of variation at this location compared with the Carrot Creek AMS and Edmonton East AMS.

The 25th, 50th, 75th, and maximum concentration values at the Edmonton East AMS were generally lower than that observed at the Genesee AMS and Carrot Creek AMS. This is consistent with central Alberta O₃ trend data from the last seven years reported by Kindzierski et al. (2006) and Kindzierski (2006). A primary reason for these lower concentrations observed at the Edmonton East AMS is due to scavenging with NO_x from urban motor vehicle emissions.

Figure 4-8 shows the cumulative distribution and key percentile values for hourly O₃ concentrations at the Genesee AMS during 2007. Hourly O₃ concentrations were <113 µg/m³ for 98 percent of time during 2007.




Figure 4-6 Cumulative Distribution Plot of 1-hour Average SO₂ Concentrations at Air Monitoring Stations in the Wabamun-Genesee Area during 2007




Figure 4-7 Box-and-Whisker Plot of 1-hour Average O₃ Concentrations at Selected Air Monitoring Stations in Central Alberta during 2007

Note: 25th %ile (bottom of box), 50th %ile (line inside box), 75th %ile (top of box), maximum (top T), and 98th %ile (diamond symbol).




Figure 4-8 Cumulative Distribution Plot of 1-hour Average O₃ Concentrations at the Genesee Air Monitoring Station during 2007

Fine Particulate Matter – Figure 4-9 is a box-and-whisker plot of 24-hour PM_{2.5} concentrations measured at the Genesee AMS and Powers AMS in 2005, 2006 and 2007. Median (50th percentile) 24-hour PM_{2.5} concentrations at both stations were low, $\leq 4 \mu\text{g}/\text{m}^3$ for all years. The 98th percentile 24-hour PM_{2.5} concentrations at both air monitoring stations were low ($\leq 14 \mu\text{g}/\text{m}^3$) for all years. These values are well below the 98th percentile, three-year average CWS of $30 \mu\text{g}/\text{m}^3$ also shown in Figure 4-9. Interquartile ranges for 24-hour PM_{2.5} concentrations at both air monitoring stations were very narrow ($\leq 5 \mu\text{g}/\text{m}^3$), indicating very little variation in 24-hour average concentrations throughout 2005, 2006 and 2007.




Figure 4-9 Box-and-Whisker Plot of 24-hour Average PM_{2.5} Concentrations at the Genesee and Powers Air Monitoring Stations (2005, 2006 and 2007)

Note: 25th %ile (bottom of box), 50th %ile (line inside box), 75th %ile (top of box), maximum (top T), and 98th %ile (diamond symbol).

It is noted that both the Genesee AMS and Powers AMS had maximum 24-hour average concentrations exceeding the three-year average CWS of 30 µg/m³ in 2006. These exceedance-days occurred in July during a high point in the Alberta forest fire season and the region was noted as being smoky at the time (TransAlta and EPCOR 2007).

Figure 4-10 shows the cumulative distribution plot for hourly PM_{2.5} concentrations measured at the Genesee and Powers air monitoring stations. Overall, hourly PM_{2.5} concentrations were low at these two stations, <16 µg/m³ for 98 percent of the time during 2007.




Figure 4-10 Cumulative Distribution Plot of 1-hour Average $\text{PM}_{2.5}$ Concentrations at the Powers and Genesee Air Monitoring Stations during 2007

4.1.3 Wind Speed and Direction Statistics

A wind rose plot is used to show information about the distribution of wind speeds, and frequency of the varying wind directions at the air monitoring stations. These plots are derived from hourly meteorological observations of wind speeds and directions. The wind rose plots shown here were divided into 16 sectors, one sector for each 22.5 degrees of the horizon. The radius of the 16 outermost, wide wedges in a wind rose plot gives the relative frequency of each of the 16 wind directions (i.e., the percent of the time that the wind is blowing from that direction). The colours indicate the wind speed range (in m/s) for a particular direction. Another common method of displaying a year of wind data at the air monitoring stations is a wind class frequency distribution plot. These plots show the percent of time that different wind speeds occur.

Figures 4-11 to 4-18 show annual wind rose plots and annual wind class frequency distribution plots for the four air monitoring stations in the Wabamun–Genesee area during 2007. Figure 4-11 shows that winds blew predominantly from the northwest and the southwest directions at the Genesee AMS (about 47 percent of the time). In the case of the Powers AMS (Figure 4-13), winds blew from the northwest and the southwest directions about 45 percent of the time.

Figure 4-15 shows that winds blew from the northwest and the southwest directions at the Meadows AMS about 47 percent of the time. In the case of the Wagner AMS (Figure 4-17), winds blew from the northwest and the southwest directions about 45 percent of the time.




Figure 4-11 Annual Wind Rose Plot for the Genesee Air Monitoring Station during 2007




Figure 4-12 Annual Wind Class Frequency Distribution Plot for the Genesee Air Monitoring Station during 2007




Figure 4-13 Annual Wind Rose Plot for the Powers Air Monitoring Station during 2007




Figure 4-14 Annual Wind Class Frequency Distribution Plot for the Powers Air Monitoring Station during 2007




Figure 4-15 Annual Wind Rose Plot for the Meadows Air Monitoring Station during 2007




Figure 4-16 Annual Wind Class Frequency Distribution Plot for the Meadows Air Monitoring Station during 2007




Figure 4-17 Annual Wind Rose Plot for the Wagner Air Monitoring Station during 2007




Figure 4-18 Annual Wind Class Frequency Distribution Plot for the Wagner Air Monitoring Station during 2007

4.2 Intermittent Monitoring Program

Results of the intermittent PM monitoring program are shown in Figure 4-19 (PM_{2.5}) and Figure 4-20 (PM₁₀). Figure 4-19 is a box-and-whisker plot of 24-hour average PM_{2.5} data collected on a 1-in-6 day schedule (NAPS). Twenty-four hour average PM_{2.5} concentrations were <20 µg/m³ at both the Genesee AMS and Powers AMS for 98 percent of the time. This is consistent with 24-hour average data observed from continuous monitoring at these stations during 2007 (Figure 4-9).




Figure 4-19 Box-and-Whisker Plot of Intermittent 24-hour Average PM_{2.5} Concentrations at Genesee and Powers Air Monitoring Stations in 2007

Note: 25th %ile (bottom of box), 50th %ile (line inside box), 75th %ile (top of box), maximum (top T), and 98th %ile (diamond symbol).

Figure 4-20 is a box-and-whisker plot of 24-hour average PM₁₀ concentrations at the Genesee AMS and Powers AMS during 2007. Twenty-four hour average PM₁₀ concentrations were <51 µg/m³ at both the Genesee and Powers air monitoring stations for 98 percent of the time. The maximum 24-hour average PM₁₀ concentration at the Genesee AMS (78 µg/m³) was higher than that at the Powers AMS (72 µg/m³). Similar variation in 24-hour average PM₁₀ concentrations was observed at Powers AMS and Genesee AMS. The IQRs were similar: 11 µg/m³ at Genesee AMS and 12 µg/m³ at Powers AMS.

Figures 4-21 and 4-22 show 24-hour average PM_{2.5} concentrations versus PM₁₀ concentrations at the Genesee and Powers air monitoring stations during 2007. Regression analysis was undertaken to examine the degree of comparison between individual 24-hour average PM_{2.5} concentrations and PM₁₀ concentrations. The purpose was to determine the ability of the PM_{2.5} concentrations to predict PM₁₀ concentrations.




Figure 4-20 Box-and-Whisker Plot of 24-hour Average PM₁₀ Concentrations at Genesee and Powers Air Monitoring Stations in 2007

Note: 25th %ile (bottom of box), 50th %ile (line inside box), 75th %ile (top of box), maximum (top T), and 98th %ile (diamond symbol).




Figure 4-21 Twenty-four Hour Average PM_{2.5} Concentration versus PM₁₀ Concentration at the Genesee Air Monitoring Station during 2007




Figure 4-22 Twenty-four Hour Average PM_{2.5} Concentration versus PM₁₀ Concentration at the Powers Air Monitoring Station during 2007

Coefficient of determination (R^2) values were calculated from PM_{2.5}/PM₁₀ matched pair data for each air monitoring station (Genesee and Powers) and are shown in Figures 4-21 and 4-22. A coefficient of determination (R^2) is the proportion of sample variance of a response variable (PM₁₀ concentration) that is "explained" by predictor variables (PM_{2.5} concentrations) when a linear regression is done.

Results for the Genesee AMS ($R^2 = 0.20$, $n=59$) indicated that only 20 percent of variance in 24-hour average PM₁₀ concentrations was explained by variance in 24-hour average PM_{2.5} concentrations. These findings indicate that 24-hour average PM_{2.5} measurements at the Genesee AMS would be poor predictors of expected 24-hour average PM₁₀ concentrations.

Results for the Powers AMS ($R^2 = 0.23$, $n=60$) indicated that only 23 percent of variance in 24-hour average PM₁₀ concentration was explained by variance in 24-hour average PM_{2.5} concentrations. Again, these findings indicate that 24-hour average PM_{2.5} measurements at Powers AMS would be poor predictors of expected 24-hour average PM₁₀ concentrations.

4.3 Passive Monitoring Program

Bubble plots are used to show spatially the size of measured values. In this case, bubble plots were used to show annual average concentrations of NO₂, SO₂, and O₃ at the 21 passive monitoring sites in the Wabamun–Genesee area during 2007 (Figure 1-2). Annual average concentrations were determined by averaging 12 monthly monitoring results at each passive site.

In addition, annual average concentrations of NO₂, SO₂, and O₃ at selected continuous monitoring stations in the area during 2007 were plotted for comparison purposes (Violet Grove AMS and Edmonton East AMS). This information was obtained from the CASA Data Warehouse for 2007 (CASA 2008).

4.3.1 Nitrogen Dioxide

Figure 4-23 is a bubble plot of annual average NO₂ concentrations measured at 18 NO₂ passive monitoring sites during 2007. Figure 4-23 indicates that annual NO₂ concentrations determined by passive monitors increase from sites near Lake Wabamun to sites adjacent to the City of Edmonton. This spatial trend is consistent with that reported by Kindzierski (2002) for the same area. This trend is due to the influence of increasing urban source NO₂ emissions within and adjacent to the City of Edmonton.

The highest annual average NO₂ concentration shown in Figure 4-23 was observed at the Edmonton East station (15.6 ppb or 29 µg/m³) compared to a range of 2.4 to 3.9 ppb (4.5 to 7.3 µg/m³) observed at passive sites located within 30 km of the generating plants. These data clearly show that urban source emissions contribute more to ambient NO₂ concentrations observed in the City of Edmonton and surrounding area relative to generating plant emissions from the Wabamun–Genesee area.

4.3.2 Sulphur Dioxide

Figure 4-24 is a bubble plot of annual average SO₂ concentrations measured at 11 SO₂ passive monitoring sites during 2007. Annual average SO₂ concentrations were very low – in the 1.0 to 2.0 ppb (2.6 to 5.2 µg/m³) range across all sites. These low values and associated lack of spatial trend is consistent with that reported by Kindzierski (2002) for the same area, based on a passive monitoring study conducted by Alberta Environment between July 2000 and September 2001 (Myrick 2002). This lack of trend is also consistent with predictions of future cumulative SO₂ emissions in the area associated with the Genesee generating plant expansion (EPCOR 2001), as well as with concentrations measured in the continuous program discussed previously in Section 4.1.2.

4.3.3 Ozone

Figure 4-25 is a bubble plot of annual average O₃ concentrations measured at 12 O₃ passive monitoring sites during 2007. In general, slightly higher annual average O₃ concentrations were observed at a location to the west (Violet Grove AMS) compared with passive monitoring sites directly east of the Wabamun–Genesee area in Figure 4-25.

A partial explanation for this spatial trend is because of O₃ scavenging which is believed to be due to increased NO_x emissions approaching the City of Edmonton (Kindzierski et al. 2006; Kindzierski 2006). This spatial trend is consistent with slightly increasing NO₂ concentrations observed towards the City of Edmonton, as shown in Figure 4-23.




Figure 4-23 Bubble Plot of Annual Average NO₂ Concentrations [ppb] at Passive Monitoring Sites and at Selected Continuous Monitoring Station Locations in Central Alberta during 2007




Figure 4-24 Bubble Plot of Annual Average SO₂ Concentrations [ppb] at Passive Monitoring Sites and at Selected Continuous Monitoring Station Locations in Central Alberta during 2007




Figure 4-25 Bubble Plot of Annual Average O₃ Concentrations [ppb] at Passive Monitoring Sites and at Selected Continuous Monitoring Station Locations in Central Alberta during 2007

O₃ and NO₂ were analyzed at passive monitoring sites 15 and 16 (Figures 4-23 and 4-25), which were commissioned in February 2006. These sites are situated downwind of a principal wind-flow direction (winds from the northwest) along an imaginary northwest line in which the generating plants are located. Based on annual average wind speeds in the area, these sites are located greater than two- to four-hours travel time downwind of the generating plants. Previous investigations by others (Gillani et al. 1998; U.S. EPA 1998b) have indicated that generating plant plume chemistry maturity and peak production capacities of anthropogenic O₃ and inorganic nitrogen species occurs between 30 and 100 km downwind of generating plants.

Passive monitoring sites 15 and 16 were established to examine whether O₃ levels are higher, lower, or similar compared to levels from O₃ sampling sites closer to the generating plants. These sites were established to help understand whether anthropogenic O₃ production may be occurring farther away. It was important to avoid siting these sampling locations near urban areas because the influence of NO_x scavenging becomes more dominant, resulting in lower O₃ levels.

Annual average O₃ concentrations at passive monitoring sites 15 and 16 were within five percent of the annual average O₃ concentration at Violet Grove AMS during 2007 (Figure 4-21). This difference is within the expected variation due to accuracy and precision of the O₃ passive monitoring device used (± 15 percent). These findings support the observation that anthropogenic O₃ production downwind of generating plant emissions is not readily apparent using passive monitors.

Figure 4-26 shows monthly average concentrations measured at O₃ passive monitoring sites during 2007. Highest monthly average O₃ concentrations observed in the network occurred in spring, peaking in March. This temporal trend is consistent with historical data reported by Kindzierski et al. (2006) and He et al. (2005) for other O₃ monitoring sites in west central Alberta. Seasonal variation is influenced by stratospheric intrusion of O₃ that has been frequently observed in previous studies in Alberta (He et al. 2005; Chaikowsky 2001; Sandhu 1999).




Figure 4-26 Monthly Average O₃ Concentration at Passive Monitoring Sites in Central Alberta during 2007

Note – refer to Figure 1-2 for location of passive monitoring sites.

5 Findings

A continuous program was conducted at the Genesee, Meadows, Wagner and Powers air monitoring stations in the Wabamun–Genesee area during 2007. Data capture rates for measured parameters at all air monitoring stations were above the 90 percent criterion required by the Alberta Environment Air Monitoring Directive. High up-times were indicative that equipment in the continuous air monitoring network was generally well-maintained. The Genesee AMS experienced a power failure and was down for five days in July, resulting in up-times of 84% for monitoring equipment during the month. The Meadows AMS also experienced a power failure and was down for 2.5 days in June, resulting in up-times of 89% for monitoring equipment during the month. Both air monitoring stations operated normally for the remainder of the year.

Overall, air quality was judged to be good at the monitoring stations during 2007, similar to that observed in 2006:

- Hourly NO₂ concentrations were very low at all four air monitoring stations (<54 µg/m³ for 98 percent of the time). The observed 10th highest 1-hour NO₂ concentrations ranged from 54 µg/m³ at Genesee AMS to 70 µg/m³ at Meadows AMS. For comparison purposes, the 10th highest 1-hour NO₂ concentration observed at the Edmonton East AMS during 2007 was 115 µg/m³. All of these concentrations fall well below the 1-hour Alberta Ambient Air Quality Objective of 400 µg/m³.
- Hourly SO₂ concentrations were very low at all four air monitoring stations (<33 µg/m³ for 98 percent of the time). The observed 10th highest 1-hour SO₂ concentrations ranged from 72 µg/m³ at Meadows AMS to 95 µg/m³ at Powers AMS. For comparison purposes, the 10th highest 1-hour SO₂ concentration observed at the Edmonton East AMS during 2007 was 56 µg/m³. All of these concentrations fall well below the 1-hour Alberta Ambient Air Quality Objective of 450 µg/m³.
- The annual average ozone (O₃) concentration at the Genesee AMS was 53 µg/m³; similar to that observed in 2006 and 2005. Hourly O₃ concentrations at the Genesee AMS were comparable to that observed at a rural air monitoring station in west central Alberta (Carrot Creek). Hourly O₃ concentrations at the Genesee AMS were <113 µg/m³ for 98 percent of the time during 2007. The 10th highest 1-hour O₃ concentration observed at Genesee AMS was 141 µg/m³. For comparison purposes, the 10th highest 1-hour O₃ concentrations observed at the Carrot Creek AMS and Edmonton East AMS were 139 and 135 µg/m³, respectively. All of these concentrations fall below the 1-hour Alberta Ambient Air Quality Objective of 160 µg/m³.
- Twenty-four hour PM_{2.5} concentrations at the Genesee and Powers AMS ranged from <1 to 20 µg/m³. Median (50th percentile) 24-hour PM_{2.5} concentrations at both stations were very low (≤4 µg/m³). The 98th percentile 24-hour PM_{2.5} concentrations at both air monitoring stations were low (<16 µg/m³).

During 2007, twenty-four hour average PM₁₀ and PM_{2.5} samples were collected at the Powers AMS and Genesee AMS according to a National Air Pollution Surveillance (NAPS) six-day cycle sampling frequency:

- Twenty-four hour average PM_{2.5} concentrations were <20 µg/m³ at both the Genesee AMS and Powers AMS for 98 percent of the time. This is consistent with 24-hour average data observed from continuous monitoring at these stations.
- Twenty-four hour average PM₁₀ concentrations were <51 µg/m³ at both the Genesee and Powers air monitoring stations for 98 percent of the time.

Passive air monitors were deployed at 21 stations in the Wabamun–Genesee area during 2007:

- At passive sites located within 30 km of the generating plants, annual average NO₂ concentrations ranged from 4.5 to 7.3 µg/m³, whereas at the Edmonton East AMS the annual average NO₂ concentration observed was 29 µg/m³. Annual NO₂ concentrations determined by passive monitors increased slightly from sites near Lake Wabamun to sites adjacent to the City of Edmonton.

This spatial trend is consistent with previous studies in the same area and is due to the influence of increasing urban NO_x emissions within and adjacent to the City of Edmonton. Urban source emissions are more important contributors to ambient NO₂ concentrations observed in the City of Edmonton and surrounding area, compared with generating plant emissions in the Wabamun–Genesee area.

- Annual average SO₂ concentrations were very low – in the 2.6 to 5.2 µg/m³ range – at all passive sites in the Wabamun–Genesee area. This lack of spatial trend is consistent with that reported in previous studies for the same area.
- Annual average O₃ concentrations observed at passive monitoring sites directly east of the Wabamun–Genesee area ranged from 23 to 29 µg/m³; whereas an annual average O₃ concentration of 29 µg/m³ was observed at a rural background air monitoring station 55 km west of the generating plants (Violet Grove AMS).
- Two passive monitoring sites are located greater than two- to four-hours travel time downwind of the generating plants along an imaginary northwest line in which the generating plants are located. Previous work by others has indicated that generating plant plume chemistry maturity and peak production capacities of anthropogenic O₃ and inorganic nitrogen species occurs between 30 and 100 km downwind of generating plants (within the range of where these sites are located).

Annual average O₃ concentrations at these two sites were within five percent of the annual average O₃ concentration at the Violet Grove AMS. This difference is within the expected accuracy and precision of the O₃ passive monitoring device used (±15 percent). These findings support the observation that anthropogenic O₃ production downwind of generating plant emissions is not readily apparent using passive monitors.

6 References

- Air Monitoring Directive (AMD) 1989. Monitoring and Reporting Procedures for Industry. Environmental Monitoring and Evaluation, Alberta Environment, Edmonton, AB. Available at: <http://environment.alberta.ca/632.html> (accessed March 2008).
- Alberta Environment (AENV). 2000. Air Quality Model Guideline. Pub. No. T/564. Alberta Environment, Edmonton, AB. 44 pp.
- Chaikowsky, C.L.A. 2001. Overview of Ground-Level Ozone Observations in Alberta; 1986-1998. Science and Technology Branch, Environmental Sciences Division, Alberta Environmental Protection, Edmonton, AB.
- Clean Air Strategic Alliance (CASA). 2006. Pollutants Monitored, Monitoring Methods, and Data Report for Hightower Ridge. CASA, Edmonton, AB. Available at: <http://www.casadata.org> (accessed March 2006).
- Clean Air Strategic Alliance (CASA). 2008. The CASA Data Warehouse, CASA, Edmonton, AB <http://www.casadata.org/> (accessed March 2008).
- EPCOR. 2001. Genesee Generating Station Phase 3. EUB/Alberta Environment Application and Supplemental Information. Environmental Impact Assessment Report. EPCOR Generation Inc. and EPCOR Power Development Corporation. Edmonton, AB. July 2001.
- Environment Canada. 2006a. Nitrogen Oxides – NO_x. Environment Canada, Ottawa, ON. Available at: http://www.ec.gc.ca/cleanair-airpur/NOx-WS489FEE7D-1_En.htm (accessed March 2008).
- Environment Canada. 2006b. Sulphur Oxides – SO_x. Environment Canada, Ottawa, ON. Available at: [http://www.ec.gc.ca/cleanair-airpur/Pollutants/Criteria_Air_Contaminants_and_Related_Pollutants/Sulphur_Oxides\(SOx\)-WSBBB2123F-1_En.htm](http://www.ec.gc.ca/cleanair-airpur/Pollutants/Criteria_Air_Contaminants_and_Related_Pollutants/Sulphur_Oxides(SOx)-WSBBB2123F-1_En.htm) (accessed March 2008).
- Faisal, K., Gamal El-Din, M., and Kindzierski, W.B. 2006. Establishment of ambient air quality trends using historical monitoring data from Edmonton and Fort McKay, Alberta. *Canadian Society for Civil Engineering 2006 Annual General Conference*, 23 to 26 May 2006, Calgary, AB. 9 pp.
- Gillani, N.V., Meagher, J.F., Valente, R.J., Imhoff, R.E., Tanner, R.L., Luria, M. 1998. Relative production of ozone and nitrates in urban and rural power plant plumes 1. Composite results based on data from 10 field measurement days. *Journal of Geophysical Research*, 103(D17): 22593-22616.
- Haque, N., Gamal El-Din, M., and Kindzierski, W.B. 2006. Analysis of ambient air quality trends at selected West Central Airshed Society stations: Tomahawk and Carrot Creek. *Canadian Society for Civil Engineering 2006 Annual General Conference*, 23 to 26 May 2006, Calgary, AB. 9 pp.

- He, Y., Selvaraj, M., Kindzierski, W.B., and Gamal El-Din, M. 2005. Investigation of Factors Affecting Ambient O₃ Concentrations in West Central Airshed Society Zone, Alberta. Report prepared for West Central Airshed Society, Drayton Valley, Alberta. 33 pp.
- Kindzierski, W.B. 2002. Recommendations for an Air Monitoring Area for Power Plants in the Genesee/Wabamun Area. Report prepared for EPCOR and TransAlta, Edmonton, AB. August 2002. 35 pp.
- Kindzierski, W.B., Gamal El-Din, M., Selvaraj, M., and He, Y. 2005. ANN modeling of O₃ at rural monitoring sites in the West Central Airshed Zone of Alberta. *Proceedings of the Air & Waste Management Association 98th Annual Conference and Exhibition*, 21 to 25 June 2005, Minneapolis, MN (11 pages).
- Kindzierski, W.B. 2006. Analysis of historical O₃ and PM_{2.5} trends from ambient air monitoring data in central Alberta. *Air Quality 2006: Piecing the Puzzle, The Fourth Canadian Workshop on Air Quality*, 13 to 16 February 2006, Banff, AB.
- Kindzierski, W.B., Gamal El-Din, M., and Haque, N. 2006. Ambient Air Quality Trends in West Central Airshed Society Zone. University of Alberta report prepared for West Central Airshed Society, Drayton Valley, AB. 70 pp.
- Myrick, B. 2002. Personal communication with W. Kindzierski. Myrick, B., Air Quality Analyst, Alberta Environment, Edmonton, AB. June 2002.
- Sandhu, H.S. 1999. Ground-Level Ozone in Alberta. Science and Technology Branch, Environmental Sciences Division, Alberta Environmental Protection, Edmonton, AB.
- TransAlta Utilities Corp. and EPCOR Utilities Inc. (TransAlta and EPCOR). 2007. TransAlta and EPCOR Ambient Air Monitoring Program, Wabamun–Genesee Area, 2006 Annual Report. TransAlta Utilities Corporation and EPCOR Utilities Inc., Edmonton, AB. June 2007.
- TransAlta Utilities Corp. and EPCOR Utilities Inc. (TransAlta and EPCOR). 2008a. TransAlta and EPCOR Acid Deposition Assessment Program, 2007 Annual Report. TransAlta Utilities Corporation and EPCOR Utilities Inc., Edmonton, AB. June 2008.
- TransAlta Utilities Corp. and EPCOR Utilities Inc. (TransAlta and EPCOR). 2008b. TransAlta and EPCOR Mercury Assessment Program, 2007 Annual Report. TransAlta Utilities Corporation and EPCOR Utilities Inc., Edmonton, AB. June 2008.
- U.S. EPA. 1998a. NO_x - How Nitrogen Oxides Affect the Way We Live and Breathe, EPA-456/F98-005. US Environmental Protection Agency, Washington, DC. Available at: <http://www.epa.gov/air/urbanair/nox/index.html> (accessed March 2008).
- U.S. EPA. 1998b. Guideline on Ozone Monitoring Site Selection. EPA-454/R-98-002. Office of Air Quality, Planning and Standards, US Environmental Protection Agency, Research Triangle Institute, NC. August 1998.
- U.S. EPA. 2000. SO₂ – How Sulfur Dioxide Affects the Way We Live and Breathe. U.S. Environmental Protection Agency, Washington, DC. Available at: <http://www.epa.gov/ebtpages/airairposulfuroxidesso2.html> (accessed March 2008).

United States Environmental Protection Agency (U.S. EPA). 2006. Particulate Matter. U.S. Environmental Protection Agency, Washington D.C. Available at: <http://www.epa.gov/air/particlepollution/index.html> (accessed March 2008).

WBK and Associates Inc. (WBK). 2007. Ambient Air Monitoring Program Quality Assurance Plan for West Central Airshed Society, Version 1.0. Report prepared for West Central Airshed Society, Drayton Valley, AB. April 2007. 81 pp.

Xu, W., Gamal El-Din, M., and Kindzierski, W.B. 2006. Trend analysis of historical ambient air monitoring data in Edmonton and Fort McKay, Alberta. *Proceedings of the Air & Waste Management Association 99th Annual Conference and Exhibition*, 18 to 22 June 2006, New Orleans, LA (15 pages).

APPENDIX Summary of 2007 Passive Monitoring Data

Table A1. Summary of 2007 NO₂ Passive Monitoring Data (monthly average concentration in ppb)

Site	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	7.8	7	6.2	4	3.1	3.3	2.5	2.9	3	5.9	5.7	10
2	4.4	6.4	4.7	3.2	2	1.8	2.4	1.7	2.8	6.8	4.4	7.3
3	5.4	5.4	4.6	2.3	1.5	1.7	1.6	1.9	2	3.7	3.4	5.7
4	4.7	5	3.6	2.5	1.7	1.4	2.6	1.5	1.5	-	3	6
4b	3.5	4.5	3.9	2.4	2.1	1.5	1.9	1.4	3	4.7	3.7	5.1
4c	6.5	6.2	4.6	2.7	2	2.2	2.3	2.1	2	4.7	3.2	8
4d	3.6	6.6	3.9	2.4	2.1	2.2	2.5	2.4	2	3.7	2.8	4.7
4e	3.4	3.8	2.9	1.7	1.4	1.4	1.6	1.4	1.6	3.1	2.7	3.8
5	3.7	4.8	3.1	2.7	2	3.1	2.7	2.2	2.7	6	3.8	5.8
6	3.4	5.4	4.2	2.3	2.2	2.8	1.8	2.2	2.7	4.2	4	5.8
7	4.9	5.9	5	3.5	3.1	3.4	5.1	2.6	3.3	5.4	4.9	7.2
8	6.7	6.9	5.2	2.7	2.4	3	2.6	1.9	3	5.3	5	8.4
9	2.1	3.9	2.8	2.3	1.9	2.6	2	1.7	2.1	3.5	2.7	3.8
10	5.9	5.4	4.1	2.4	2.1	2.8	2.4	2.2	2.9	5.4	4.5	6.1
11	5	5.1	4.1	2.1	1.9	3	3.2	2	3.3	4.2	3.7	2.9
15	3.5	4.3	3.8	2.1	1.7	1.8	1.9	2	2.3	3.6	3.8	4.2
16	4.5	4.7	4.7	2.2	1.8	2.3	1.9	2.2	2.7	4.2	3.7	2.8
Genesee	3.8	4.5	3.5	2.3	2.5	2.5	2.4	3.7	2.9	4.1	3.5	4.8

- not available

Table A2. Summary of 2007 SO₂ Passive Monitoring Data (monthly average concentration in ppb)

Site	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
3	0.6	2	1.5	0.7	0.9	1	1.1	0.6	0.5	1.2	0.5	0.8
4	0.9	2.1	1.5	1	0.9	0.9	1.4	1	0.7	-	0.4	0.8
4b	1	3	2.2	1.3	1.6	0.8	1.7	0.9	0.8	2.3	1	1.8
4c	0.9	3.1	2.1	0.8	0.9	0.8	1.5	1.5	0.4	1	0.6	1.2
4d	1	2.6	1.6	0.9	1.2	1.6	2.3	1.8	0.8	1.4	0.5	1.3
4e	1.1	2.2	1.7	0.8	0.8	1	1.2	1.3	0.6	1.3	0.7	1.2
5	1.3	2	1.6	1.1	1.5	1.4	2.2	0.8	1.2	2.8	1.3	1.2
6	1.3	2.5	1.6	0.9	0.8	1	1.1	1.1	0.8	1.9	0.9	1.1
9	0.8	2.1	2.1	1.1	1.5	1.5	1.6	2	1.2	1.7	0.6	1.3
10	1.7	2.7	1.8	0.7	1.1	1.3	1.7	1.8	1.4	1.6	1.4	1.4
Genesee	1.5	2.5	2.1	1.4	2	1.6	2.3	3.2	2.3	1.6	1.5	1.8

- not available

Table A3. Summary of 2007 O₃ Passive Monitoring Data (monthly average concentration in ppb).

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	26.7	30.6	38.6	35	33.6	25.2	22.7	17.6	12	17.3	20.1	23.8
3	22.5	28.4	36	30.5	27.4	24.1	21.2	15.3	14.2	16.4	19	19.9
6	29.2	34	35.4	34.9	33.1	27.4	25.4	18.6	19.5	20.8	21.8	25.6
7	30.2	34.2	39.4	35.5	33.6	26.2	26.7	17.8	19	20.4	22	25.1
10	30	33.9	39.3	35.4	31.9	25.4	25.6	16.6	21.2	20.7	19.4	25.9
11	31.6	39.6	39.9	37	29.6	26.3	23.3	16.4	17.9	18.6	19.6	26.9
12	25.9	30.5	31.3	32.3	28.6	26.6	23.5	16.8	23.8	17.8	17.2	19.9
13	13.5	19.5	23.2	30.9	29.1	27	31.4	16.8	18.3	16	13.9	10.8
14	20.7	28.1	29.7	31.9	28.2	28.3	27.5	20	16.4	16.3	17.5	16.5
15	36.2	39.6	41.4	38.6	32.7	27.4	27.9	19.7	19.1	24.4	24.5	31.8
16	32.1	38.8	38.4	40.3	32	28.9	30.9	22.2	23.7	23.4	26	30.2
Genesee	29.5	37.6	41.6	37.8	31.5	25.6	28.1	19.4	18	21.3	22.6	29.9